

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Report from Seattle, Wash.—Plague-prevention work.

Passed Assistant Surgeon Glover reports November 2:

SEATTLE, WASH.

Week ended October 31.

Date of finding of last plague rat, September 26, 1908.	
Rats received	1,310
Rats necropsied	1, 180
Plague-infected rats to date	21
Vessels inspected.	12
Vessel fumigated	

STATISTICAL REPORTS OF MORBIDITY AND MORTALITY, STATES AND CITIES OF THE UNITED STATES—UNTABULATED.

CONNECTICUT—Stamford.—Month of October, 1908. Estimated population, 22,000. Number of deaths not reported. Cases: Enteric fever 1, scarlet fever 2, tuberculosis 2.

FLORIDA.—Reports from the state board of health for the week ended October 31, 1908, show as follows: Enteric fever—Tampa, 1 case. Tuberculosis—Jacksonville, $Lake\ City$, and Ocala, each 1 case. Diphtheria—Bradentown, 3; Jacksonville, and Tampa, each 2 cases; $Ybor\ City$, Pensacola, Palmetto, and Arcadia, each 1 case.

Pensacola.—Month of October, 1908. Estimated population, 30,000. Total number of deaths, 28, including enteric fever 2 and 1 from tuberculosis.

Indiana—Muncie.—Month ended November 4, 1908. Estimated population, 34,036. Total number of deaths, 18, including 4 from tuberculosis. Cases: Enteric fever 6, scarlet fever 13, diphtheria 3, tuberculosis 3.

Iowa—Cedar Rapids.—Month of October, 1908. Estimated population, 30,000. Total number of deaths, 34, including enteric fever 2 and 1 from tuberculosis. Cases: Scarlet fever, 4.

Kentucky—Henderson.—Four months ended October 31, 1908. Estimated population, 17,500. Total number of deaths, 75, including enteric fever 2 and 4 from tuberculosis. Cases: Enteric fever 17, scarlet fever 3, measles 1, whooping cough 6, and 5 from tuberculosis.

NEW HAMPSHIRE—Concord.—Month of October, 1908. Estimated population, 20,000. Total number of deaths, 33, including 3 from tuberculosis. Cases: Enteric fever 3, diphtheria 4, and tuberculosis 3.

Franklin.—Month of October, 1908. Estimated population, 7,200. Total number of deaths, 9, including 1 from tuberculosis. Cases: Tuberculosis 1 and enteric fever 1.

West Virginia—Charleston.—Month of October, 1908. Estimated population, 26,000. Total number of deaths, 17, including enteric fever 1, diphtheria 11, and 1 from tuberculosis. Cases: Diphtheria 9, scarlet fever 3, enteric fever 7, and tuberculosis 1.